

- Sodium thiosulfate
- Potassium iodate
- Potassium iodide
- Starch solution (freshly made)
- Sulfuric acid, 1 mol dm³
- Sample(s) of fruit juice

Stock solutions

1. Sodium thiosulfate solution 0.010 mol dm⁻³
Weigh out, accurately, ca 0.620 g of Na₂S₂O₃·5H₂O, dissolve in deionised water and make up to 250 cm³ in a volumetric flask. Store this stock solution in a dark glass bottle.
2. Potassium iodate solution 0.001 mol dm⁻³
Weigh out, accurately, ca 0.054 g of KIO₃, dissolve in deionised water and make up to 250 cm³ in a volumetric flask.
3. Potassium iodide solution 0.005 mol dm⁻³
Weigh out 0.21 g of KI, dissolve in deionised water and make up to 250 cm³ with deionised water.

Observations

The titre volume should be in the range 0.5–1 cm³, the disappearance of the blue-black colour marking the end-point.

This experiment offers possibilities for assessing students' abilities in following instructions and/or processing results.

A survey of a range of fruit drinks (and maybe other products containing vitamin C) could form the basis of a class project or as an activity for a school or college chemistry club.

Notes

The reaction to generate the iodine is based on using an accurately known volume of the potassium iodate solution (the concentration of which is accurately known).

The potassium iodide solution and the sulfuric acid are added in slight excess, and thus the concentrations of these solutions is not critical.

Instead of generating the iodine in situ, it is possible to use standard iodine solution in this procedure.

This would need to be diluted to give an aliquot containing 7.2×10^{-6} moles of iodine for each determination.

Specimen result and calculation

Volume of thiosulfate delivered during the titration = 0.74 cm³.

Concentration of thiosulfate = 0.010 mol dm⁻³.

Therefore, number of moles thiosulfate =

$$\frac{0.74 \times 0.01}{1000} = 7.4 \times 10^{-6}$$

Therefore, the number of moles of iodine that this reacted with during the titration = 3.7×10^{-6} .

The total number of moles of iodine produced in the reaction between iodate, iodine and sulfuric acid based on using 2 cm^3 of iodate with a concentration of $0.0012 \text{ mol dm}^{-3}$ =

$$\frac{3 \times 2 \times 0.0012}{1000} = 7.2 \times 10^{-6}$$

Therefore, the number of moles of iodine which reacted with the ascorbic acid = $7.2 \times 10^{-6} - 3.7 \times 10^{-6} = 3.5 \times 10^{-6}$

Since 1 mole of iodine reacts with 1 mole of ascorbic acid then the number of moles of ascorbic acid is also 3.5×10^{-6} .

The volume of the fruit juice used was 1 cm^3 .

Therefore, the number of moles of ascorbic acid in $1000 \text{ cm}^3 = 3.5 \times 10^{-3}$.

The relative molar mass of ascorbic acid = 174.12 g .

Therefore, mass of ascorbic acid (in 1000 cm^3) = $174.12 \text{ g} \times 3.5 \times 10^{-3} = 0.609 \text{ g}$.

The vitamin C content of the fruit drink = $61 \text{ mg per } 100 \text{ cm}^3$.

Health, safety and technical notes

- Read our standard health and safety guidance here <https://rsc.li/3SRKCow>
- Wear eye protection.
- Sulfuric acid, 1 mol dm^{-3} is a skin/eye irritant (see CLEAPSS Hazcard [HC098a](#)).
- Sodium thiosulfate, $0.010 \text{ mol dm}^{-3}$, potassium iodate, $0.001 \text{ mol dm}^{-3}$ and potassium iodide, $0.005 \text{ mol dm}^{-3}$ solutions are of low hazard, as are the starch solution and fruit juices (see CLEAPSS Hazcards [HC095a](#), [HC080](#), [HC047b](#)).